

## **Mutation Induction in Physic Nut (*Jatropha curcas* L.)**

### **by Colchicine Treatments**

Pantipa Na Chiangmai\*, Yupa Pootaeng-on, Pimjai Meetum, Neeranoot Jankomon,  
Daungruethai Muangnoi and Duanpen Kitthip

*Faculty of Animal Sciences and Agricultural Technology, Silpakorn University,  
Phetchaburi IT Campus, Cha-Am, Phetchaburi, Thailand*

*\*Corresponding author. E-mail address: pantipa@su.ac.th, mchiangmai@gmail.com*

#### **Abstract**

The objective of this study aimed to evaluate the genetic variation in physic nut (*Jatropha curcas* L.) by colchicine induction in two experiments. In Experiment 1, inflorescences of physic nut were treated under different colchicine concentrations by dropping the solution directly at the stage that the frequency of male flower at microspore stage. In Experiment 2, seeds were soaked in different concentrations of colchicine at various durations.

In Experiment 1 and 2, the results showed that the increasing of colchicine concentration likely to decreased size of guard cells, plant height and stomatal density. However, significant different among those parameters was not observed. The ploidy level determined from leaf cells of plants by flow cytometry in both experiments showed almost of plants were disomic (2x). However, in experiment 1, incomplete aneuploidy or mixoploid plants that mixing between trisomic ( $2x + 1$ ) (at 0 mM colchicine) and tetrasomic ( $2x + 2$ ) (at 2.5 mM colchicine) with disomic were also found. In experiment 2, incomplete aneuploid plants were composed of trisomic, tetrasomic and pentasomic ( $2x + 3$ ) with disomic. The mixoploid plant of disomic, trisomic and hexasomic ( $2x + 4$ ) was also detected when concentrations of colchicine and times for soaking duration were increased.

**Key Words:** Aneuploid; Energy plant; Genetic variability; Mixoploid; Ploidy chromosome level

#### **Introduction**

The issue of looming energy crisis around the world (Kung and Tam, 2005), including Thailand, has prompted researchers to pay more attention to research in an alternative energy resources (Paepatung et al., 2009; Peerapong and Limmeechokchai, 2009). Physic nut (*Jatropha curcas* L.), an energy plant that has high oil content in seed, is one of the prime target under investigation by various researchers (Heller, 1996; Trabi et al.,

1997; Winkler et al., 1997). Apart from its high oil content, physic nut is also capable of growing in infertile soil (Heller, 1996; Ginting and Maryono, 2009), making it one of the ideal trees which should be promoted for planting to provide oil for domestic use.

However, one of its shortcoming trait is that these seeds with high oil yield do not reach maturity synchronously, contributing to a low yield of fresh nut per plant during harvesting and requiring many

trees to obtain enough matured seeds in a single harvest for oil extraction (Jongschaap et al., 2007; Sharma et al., 2013). Improving physic nut using conventional breeding may encounter difficulty as this plant has narrow genetic diversity (Basha and Sujatha, 2007; Ambrosi et al., 2010; Songsri et al., 2011).

To solve these problems, the genetic improvement by inducing mutation through the callus of physic nut has been investigated by various researchers (Sujatha and Mukta, 1996; Sardana et al., 2000; Rajore and Batra, 2005; Sujatha et al., 2006; Wei et al., 2004). Tetraploid was successfully induced from diploid chromosome set of physic nut using colchicine (Sujatha and Prabhakaran, 2003; Piromya and Kermanee, 2013). However, proper protocol in mass propagation after obtaining a mutant was required and tissue culture, as one of the propagation technique, had been employed to regenerate this mutant tissue into new plants (Sujatha and Sailaja, 2005; Li et al., 2006). The dominant method for mutant induction in plant to be polyploidy was established either *in vitro* only or combined using between *in vitro* and *ex vitro* (Carretero et al., 2007; Dasgupta et al., 2008; Oates et al., 2013). This study aimed to investigate the effect of colchicines on inducing mutant in both inflorescence at microspore stage and seed embryos of physic nut by soaking.

## Materials and Methods

### Plant genetic and site of plantation

In experiment 1, the effect of treating inflorescences of a 1-year-old physic nut trees; *Jatropha curcas* (Phetchaburi variety) with colchicines was investigated. The trees were planted using the spacing 2 x 2 m<sup>2</sup>. This experiment was carried out from July to October, 2010, at Phetchaburi, Thailand. The site of plantation has an elevation approximately 0.4 meter above sea level at N12°

37.813 ´ E099° 51.588 ´ . The study was conducted in rainy season with an average rainfall of 136.6 mm in July and 212.7 in October. The averages of temperature were 27.5 °C in July and 27.8 °C in October. The averages of duration of sunshine were 4.6 hours/day in July and 4.0 hours/day in October. The field facility was located at Agricultural Farm of the Phetchaburi College of Agriculture and Technology, Cha-am District, Phetchaburi Province, Thailand.

In experiment 2, the effect of soaking seeds of physic nut trees with colchicines on 0, 0.25, 1.25 and 2.50 mM was studied. Seeds of *J. curcas* (Phetchaburi variety) harvested from the physic trees (in the plantation as described above) were used in this experiment. This experiment was carried out from August to October, 2010 in the green house. The seeds from either treated inflorescences (from Experiment 1) or treated seeds (from Experiment 2) were planted in the green house and many characteristics were determined. These included the ratio of female and male flowers after treating by different colchicine concentrations in the inflorescences in Experiment 1. The characteristics of the germinated seedlings (both from treated inflorescences and treated seeds) including plant height, stomatal density and the sizes of guard cell were determined in the laboratory at Animal Sciences and Agricultural Technology, Silpakorn University, Information Technology Campus, Cha-am District, Phetchaburi Province, Thailand. Ploidy level was determined by flow cytometry at Plant Biotechnology Laboratory, Kasetsart University, Kamphaeng Saen Campus, Kamphaeng Saen District, Nakhon Pathom Province, Thailand.

### Induce mutation

In experiment 1, five inflorescences of physic nut tree were collected from each tree and they were treated with colchicines at different concentrations (at 0, 2.5, 5, 7.5 and 10 mM). This was carried

out by dropping this chemical agent on the cotton sheet holding an individual inflorescence, with seven replicates for each colchicine concentration. The colchicine was applied at 6.30 am for 7 days consecutively.

In experiment 2, one hundred and thirty seeds of physic nut, variety Pechaburi were used in this experiment. Seeds were soaked with colchicines at different concentrations (0, 0.25, 1.25, 2.50 mM colchicine). These seeds were soaked for 6, 12, 24 and 48 hours for each concentration with 10 replications.

#### **Characteristic determination and ploidy level determination**

In the green house, height of the plantlet (obtained from either treated inflorescence or seed with different concentrations of colchicine at various durations) was measured 14 days after transplanting to the pot (in the Experiment 1 and 2, respectively). At 60 days after planting, young leaves (leaf no. 2-3 from apical shoot) in each plant were determined with respect to their stomatal density, width and length of guard cells, (with microscope). The ploidy level determined with flow cytometry (Partec ploidy Analyse PA II). The assay procedure for preparing material was carried out using 50 mg of plant tissue. The tissue was placed a plastic petri dish, added 0.5 ml of CyStain UV ploidy and chopped the tissue with a sharp razor blade for releasing the nuclei from the cells. Then, added 1.5 ml of CyStain UV ploidy and incubate at room temperature for 5 minutes. The sample was filtered through a Partec 50  $\mu$ m CellTrics disposable filter and to analyzed in flow cytometer using UV excitation (HBO-lamp, UV-laser) and measure blue emission.

#### **Experimental design and statistical analysis**

Randomized completed block design (RCBD) and completely randomized design (CRD) were conducted in the experiment 1 and 2, respectively.

Analysis of variance (ANOVA) was conducted and means among treatment were compared with the Duncan's multiple range test (DMRT).

### **Results**

#### **Experiment 1: Effect of colchicine on morphological and physiological characters of physic nut under field condition.**

The results were shown in Table 1 and 2. Non-significant different was observed on ratio of female and male flowers of physic nut (ranged from 0.043-0.117) after the inflorescences were treated with different colchicine concentrations (Table 1).

Different concentration of colchicine did not affect plant height at 14 days after planting the seedling (from treated inflorescences), with the values ranging from 9.50-11.37 cm (Table 1).

At 60 days after planting of physic nut, the density of guard cells from young leaf was not significantly affected by concentrations of colchicine, in which the density of guard cells ranged between 9.36-10.42 guard cells  $\text{mm}^{-2}$ . The highest stomata density was found in the nil control treatment (0 mM) at 10.42 guard cells  $\text{mm}^{-2}$  (Table 1).

Both width and length of guard cells of the young leaf were significantly affected by colchicine concentrations (Table 1). The width of guard cell was the highest at 0 mM colchicine (0.410 mm) and decreased when higher concentrations of colchicine were applied. The lowest width at 0.030 mm was obtained from treating with colchicines at concentration of 7.5 mM.

For the length of guard cell, the longest of them was found in seedling from treating inflorescence with 5 mM colchicine at 0.040 mm while the lowest length was detected at 2.5 mM (0.034 mm) and 7.5 mM (0.033 mm) colchicine.

The result of ploidy level from leaf of physic nut detected by flow cytometry was shown in Table

**Table 1** Characteristics of female/male flowers ratio, plant height, stomatal density and guard cell size from leaf of physic nut seedling (mean  $\pm$  SE) plants obtained from fertilized ovule with colchicine-treated inflorescences (Experiment 1)

Colchicine concentrations (mM)	Female/male flowers ratio	Plant height (cm) (at 14 DAS <sup>†</sup> )	Stomatal density (per mm <sup>2</sup> ) (at 60 DAS)	Width of guard cell (mm) (at 60 DAS)	Length of guard cell (mm) (at 60 DAS)
0	0.081 $\pm$ 5.70	10.68 $\pm$ 3.58	10.42 $\pm$ 0.28	0.410 $\pm$ 8x10 <sup>-5</sup> a	0.039 $\pm$ 4.3x10 <sup>-3</sup> ab
2.5	0.058 $\pm$ 6.64	10.43 $\pm$ 1.54	9.95 $\pm$ 0.38	0.034 $\pm$ 1.2x10 <sup>-4</sup> ab	0.034 $\pm$ 1.6x10 <sup>-3</sup> c
5	0.177 $\pm$ 0.08	9.85 $\pm$ 1.20	9.36 $\pm$ 0.27	0.038 $\pm$ 2.0x10 <sup>-4</sup> ab	0.040 $\pm$ 0.00 a
7.5	0.043 $\pm$ 6.01	11.37 $\pm$ 0.37	9.48 $\pm$ 0.12	0.030 $\pm$ 0.00 c	0.033 $\pm$ 7x10 <sup>-4</sup> c
10	0.049 $\pm$ 5.70	9.50 $\pm$ 2.12	9.71 $\pm$ 0.07	0.036 $\pm$ 2.0x10 <sup>-4</sup> ab	0.035 $\pm$ 6.0x10 <sup>-4</sup> bc
F-test	Ns <sup>‡</sup>	Ns	Ns	**	*

<sup>†</sup> DAS, days after seeding.

<sup>‡</sup> Ns, not significant difference at the 0.05 level of probability.

\* significant difference at the 0.05 level of probability.

\*\* significant difference at the 0.01 level of probability.

2. From 22 plants (plant no. 40 to no. 61), disomic (2x) was found in 19 plants (86.36%). Three (13.64%) possessed mixoploid which mixed DNA concentrations (2 peaks), trisomic (2x + 1)(0 mM colchicine) (plant no. 40), tetrasomic (2x + 2)(0 and 2.5 mM colchicine) (plant no. 41 and 42) and disomic (Table 2).

#### **Experiment 2: Effect of colchicine on growth of physic nut by seeds soaking.**

The plant height at 14 days after planting was not significantly affected by the combination of colchicine concentrations and soaking durations (ranging from 7.50-11 cm) (Table 3). At 60 days after planting, stomatal density was not significant different among treatments, which the values ranged from 6.23 to 13.48 stomata mm<sup>-2</sup> (Table 3). However, size of guard cells (width and length) was significant affected by treatments, in which the width values ranged from 0.027 to 0.046 mm and the length values ranged from 0.037 to 0.047 mm (Table 3).

Flow cytometry study showed that the

plantlets possessed both disomic and mixoploid plants (Table 4). Mixoploid of trisomic (plant no. 16), tetrasomic (plant no. 21, 23, 26, 31 and 34) and pentasomic (2x + 3) (plant no. 2) were observed. Moreover, the mixoploid plant of disomic, trisomic and hexasomic (2x + 4) also was found at 2.5 mM colchicine after soaking the seeds for 6 hrs (plant no. 3).

#### **Discussion**

##### **Experiment 1: Effect of colchicine on morphological and physiological characters of physic nut under field condition.**

The ratio of female and male flowers of physic nut (0.043-0.177) was not affected by colchicine concentrations. However, the ratio was decreased at high concentration (Table 1). The decrement of female and male flowers in the same inflorescence occurred after treatment with different concentrations of colchicine. Increase in concentrations of colchicine reduced male flowers greater than female flowers.

Plant height was used as an indicator to predict the ploidy level in plant (Miller et al., 2012). Other characteristics such as stomatal size, flower size, seed weight, plant dry matter, chlorophyll and starch contents, and strength of growth were also important characteristics to evaluate the ploidy level in plant (Miller et al., 2012). Nevertheless, in this study plant height was not significantly different (at 14 days after planting) after inflorescences were treated with different concentrations of colchicine.

In this study, guard cell sizes were significantly different but the stomatal density was not affected by colchicine treatments (Table 1). The frequency of stomatal density and epidermal cell was reported to decrease and the size of guard cell increased in brome grass (Tan and Dunn, 1973), triticale (Sapra et al., 1975), *Coffea canephora* (cultivar S. 274) (Mishra, 1997) and *Arabidopsis thaliana* (Miller et al., 2012). These results may be due to both larger epidermal cells and lower density of guard cells indicating ploidy level of plants in comparison with diploid plant (Mishra, 1997). However, in this study, the stomatal sizes tended to decrease with the increase in colchicine concentration (Table 1). There are probably at least two important factors affecting both stomatal density and size of guard cells. Those are genotype and environment (such as light, temperature, moisture affected to epidermal cell size) (Fernandez and Muzica, 1973; Mishra, 1997). Thus, the determination of ploidy level using the flow cytometry was employed to enumerate ploidy in physic nut (Heslop-Harrison, 1995; Inagaki, 2003). Flow cytometry was used for estimating DNA quantity in cell nuclei, screening for ploidy, and detecting aneuploidy and mixoploid (Doležel and Bartoš, 2005).

Flow cytometry study showed that there was the mosaic of different chromosome sets or mixoploid in plant cells of the three plant samples (Table 2). Mixoploidy was the multiple of chromosome number

or the imbalance of chromosome number in cell (not complete aneuploidy). This multiple copies of chromosomes and genes located on that chromosome may affect gene expression or phenotype, in which the magnitude of an effect depended upon plant species, characteristics, and type of aneuploidy (Lee et al., 1996; Birchler et al., 2001; Birchler, 2010; Henry et al., 2010). Mixoploidy could occur by treating many plant tissues (such as seed, buds and shoots) with colchicine in different plant species (such as ryegrass, fenugreek (*Trigonella foenum-graecum* L.) and *Dioscorea zingiberensis*) (Hill and Myers, 1944; Huang et al., 2010; Omezzine et al., 2012).

Many researchers reported about the type of ploidy; trisomic (Koh et al., 1993; Henry et al., 2010) or tetrasomic (Moody et al., 1993; Jenczewski et al., 2002) in plant tissues after they were treated with chemical agents. The complex of mixoploid (the combination of disomic, trisomic and tetrasomic) was observed in the physic nut cells in this study. Tetrasomic may occur as a result of spontaneous or induced mutation (Mayo, 1971; Bever and Felber, 1991; Moody et al., 1993). The combination of many types of ploidy level may cause the complex of phenotype expression in many traits. Trisomic was reported to alter the expression of many genes, including dosage-sensitive, in phenophytes (Henry et al., 2010).

The results in this study showed mixoploidy of trisomic (No. 40) and tetrasomic (No. 41) with disomic in leaf cells from non-treated control seeds (Table 2). However, trisomic and tetrasomic had less areas than that of disomic areas in plant cells after determination with a flow cytometry.

It was possible that the progeny seeds were from the fertilization between female flower with normal monoploid chromosome (receiving 0 mM colchicine treatment) and male flower with imbalance chromosome number (extra 1;  $x + 1$  or

**Table 2** Ploidy level of physic nut prepared from young leaf of plant obtained from fertilized ovule with colchicine-treated inflorescences (Experiment 1)

Plant No.	Peak	Index	Mean	Area	Area (%)	C.V. (%)	Chi-square
40	1	1.000	210.30	1135	69.51	5.81	0.56
	2	1.406	295.72	498	30.49	43.17	0.56
41	1	1.000	197.18	1325	82.37	5.12	0.51
	2	1.981	390.56	284	17.63	4.27	0.51
42	1	1.000	178.84	1387	98.01	6.88	0.53
	2	2.281	407.88	28	1.99	0.49	0.53
43	1	1.000	180.39	1378	100.00	15.84	0.62
44	1	1.000	187.37	1349	100.00	6.40	0.61
45	1	1.000	188.69	1208	100.00	9.61	0.66
46	1	1.000	176.43	1438	100.00	15.47	0.63
47	1	1.000	172.03	1551	100.00	11.62	0.53
48	1	1.000	182.72	1324	100.00	11.20	0.60
49	1	1.000	175.80	1473	100.00	11.85	0.61
50	1	1.000	177.24	1477	100.00	13.70	0.59
51	1	1.000	176.95	16.73	100.00	8.57	0.45
52	1	1.000	169.72	1527	100.00	13.73	0.53
53	1	1.000	175.45	1461	100.00	13.76	0.57
54	1	1.000	168.77	1688	100.00	12.35	0.50
55	1	1.000	153.37	1414	100.00	9.88	0.73
56	1	1.000	177.06	1587	100.00	11.94	0.53
57	1	1.000	180.88	1771	100.00	8.19	0.45
58	1	1.000	173.27	1586	100.00	13.53	0.57
59	1	1.000	183.85	1662	100.00	14.93	0.78
60	1	1.000	180.18	1783	100.00	14.01	0.55
61	1	1.000	185.54	1527	100.00	17.44	0.82

**Table 3** Characteristics of physic nut (mean  $\pm$  SE) seedlings obtained from soaking seeds with different times and concentrations of colchicine (Experiment 2)

Colchicine concentration-Soaking duration	Plant height (cm) (at 14 DAS <sup>†</sup> )	Stomatal density (no. mm <sup>-2</sup> ) (at 60 DAS)	Width of guard cell (mm) (at 60 DAS)	Length of guard cell (mm) (at 60 DAS)
0 mM - 6 hrs. (T1)	10.42 $\pm$ 1.80	10.79 $\pm$ 0.45	0.027 $\pm$ 1x10 <sup>-4</sup> c	0.038 $\pm$ 1x10 <sup>-5</sup> ab
0 mM - 12 hrs. (T2)	8.00 $\pm$ 0.00	8.22 $\pm$ 0.00	0.042 $\pm$ 0.00 ab	0.041 $\pm$ 0.00 ab
0 mM - 24 hrs. (T3)	8.00 $\pm$ 0.00	6.23 $\pm$ 0.00	0.046 $\pm$ 0.00 a	0.044 $\pm$ 0.00 ab
0.25 mM - 6 hrs. (T4)	10.13 $\pm$ 0.85	11.14 $\pm$ 0.40	0.040 $\pm$ 6x10 <sup>-5</sup> ab	0.042 $\pm$ 8x10 <sup>-5</sup> ab
0.25 mM - 12 hrs. (T5)	9.50 $\pm$ 0.00	10.15 $\pm$ 0.00	0.042 $\pm$ 0.00 ab	0.047 $\pm$ 0.00 a
0.25 mM - 48 hrs. (T6)	9.75 $\pm$ 1.06	13.19 $\pm$ 0.76	0.038 $\pm$ 1.5x10 <sup>-4</sup> ab	0.039 $\pm$ 2.2x10 <sup>-4</sup> ab
1.25 mM - 6 hrs. (T7)	9.00 $\pm$ 1.56	13.28 $\pm$ 5.51	0.037 $\pm$ 5x10 <sup>-5</sup> b	0.037 $\pm$ 4x10 <sup>-5</sup> b
1.25 mM - 12 hrs. (T8)	8.33 $\pm$ 1.53	13.48 $\pm$ 0.63	0.036 $\pm$ 2x10 <sup>-5</sup> bc	0.042 $\pm$ 1x10 <sup>-4</sup> ab
1.25 mM - 24 hrs. (T9)	7.50 $\pm$ 3.28	13.06 $\pm$ 0.18	0.036 $\pm$ 2x10 <sup>-5</sup> bc	0.037 $\pm$ 4x10 <sup>-5</sup> b
2.50 mM - 6 hrs. (T10)	11.00 $\pm$ 1.00	11.69 $\pm$ 0.77	0.040 $\pm$ 1.5x10 <sup>-4</sup> ab	0.039 $\pm$ 6x10 <sup>-5</sup> ab
2.50 mM - 12 hrs. (T11)	8.25 $\pm$ 1.92	11.09 $\pm$ 0.48	0.038 $\pm$ 9x10 <sup>-5</sup> ab	0.043 $\pm$ 5x10 <sup>-5</sup> ab
2.50 mM - 24 hrs. (T12)	10.50 $\pm$ 2.12	12.73 $\pm$ 0.25	0.039 $\pm$ 3x10 <sup>-5</sup> ab	0.037 $\pm$ 1x10 <sup>-5</sup> b
2.50 mM - 48 hrs. (T13)	9.25 $\pm$ 0.35	7.46 $\pm$ 0.24	0.042 $\pm$ 1.2x10 <sup>-4</sup> ab	0.041 $\pm$ 8x10 <sup>-5</sup> ab
F-test	Ns <sup>‡</sup>	Ns	**	*

<sup>†</sup> DAS, days after seeding.

<sup>‡</sup> Ns, not significant difference at the 0.05 level of probability.

\* significant difference at the 0.05 level of probability.

\*\* significant difference at the 0.01 level of probability.

extra 2; x + 2 in some chromosomes) (receiving colchicine treatment) from different inflorescences. In this study, the treated inflorescences were not covered with the bag, leading to a possibility of cross-fertilization between monoploid chromosome number of female gamete and imbalance chromosome number (extra 1; x + 1 or extra 2; x + 2 in some chromosomes) of male gamete.

#### **Experiment 2: Effect of colchicine on growth of physic nut by seeds soaking.**

In this experiment, the result on plant height

was similar to that observed in Experiment 1. This trait was not significantly affected at 14 days after planting (Table 3). However, size of guard cells was affected by the concentrations of colchicine and soaking duration (Table 3). The values of width and length of the guard cells decreased when the leaves were treated with the increased concentrations of colchicines, the result similar to that in Experiment 1. DNA content quantification by flow cytometry revealed that most of the non-treated seeds had normal diploidy (Table 4), in which many types

**Table 4** Ploidy level of physic nut prepared from young leaf of plant grew from seeds soaking by colchicine treatments (Experiment 2)

Plant No.	Peak	Index	Mean	Area	Area (%)	C.V. (%)	Chi-square
1	1	1.000	200.79	1.422	100.00	9.60	0.56
2	1	1.00	247.81	1285	87.78	5.19	0.54
	2	1.977	490.00	179	12.22	4.64	0.54
3	1	1.000	237.29	578	38.79	29.60	0.58
	2	1.180	279.92	731	49.06	5.08	0.58
	3	2.395	568.30	181	12.15	4.13	0.58
5	1	1.000	186.30	1471	100.00	11.64	0.64
6	1	1.000	220.12	1357	100.00	22.47	0.70
8	1	1.000	217.87	1317	88.14	24.76	0.61
	2	1.129	246.02	177	11.86	2.78	0.61
9	1	1.000	226.26	1225	100.00	16.20	0.62
10	1	1.000	264.66	1394	100.00	7.98	0.92
11	1	1.000	248.19	1101	100.00	5.65	0.73
12	1	1.000	206.82	1136	100.00	23.67	0.63
13	1	1.000	242.37	1163	100.00	4.88	0.69
14	1	1.000	189.12	1193	100.00	9.46	0.66
15	1	1.000	184.29	1270	100.00	8.39	0.61
16	1	1.000	181.16	1327	73.13	6.86	0.42
	2	1.726	312.64	488	26.87	26.33	0.42
17	1	1.000	194.17	1563	100.00	6.52	0.46
18	1	1.000	191.40	1460	100.00	5.74	0.54
19	1	1.000	179.77	1393	100.00	15.22	0.57
20	1	1.000	186.59	1372	100.00	14.24	0.63
21	1	1.000	197.75	1479	85.81	4.70	0.41
	2	1.992	393.90	245	14.19	3.59	0.41
22	1	1.000	186.77	1459	100.00	8.19	0.67
23	1	1.000	193.56	1223	73.16	6.10	0.51
	2	2.017	390.37	449	26.84	7.33	0.51
24	1	1.000	171.94	1308	100.00	12.93	0.64
25	1	1.000	175.20	1427	100.00	6.41	0.50
26	1	1.000	195.11	1394	86.68	4.76	0.47
	2	1.970	384.35	214	13.32	5.97	0.47
27	1	1.000	201.44	1454	100.00	8.89	0.63
28	1	1.000	176.83	1692	100.00	14.93	0.91
29	1	1.000	180.13	1405	100.00	15.82	0.58
30	1	1.000	172.23	1374	100.00	6.32	0.57
31	1	1.000	195.70	1367	76.73	5.76	0.40
	2	1.976	386.75	415	23.27	4.90	0.40
32	1	1.000	183.10	1585	100.00	8.44	0.50
33	1	1.000	188.4	1387	100.00	6.26	0.67
34	1	1.000	193.69	1556	84.61	8.51	0.45
	2	1.942	376.14	283	15.39	6.63	0.45
35	1	1.000	195.41	924	100.00	11.83	0.64
36	1	1.000	185.97	1179	100.00	8.24	0.64
37	1	1.000	194.96	1217	100.00	6.50	0.63
39	1	1.000	177.02	1476	100.00	15.11	0.58

of mixoploidy were also observed in different combinations of colchicine concentration and soaking duration. The mixoploid of plant came from treated seed may cause by chimera mutant of cells composed within embryo. Using seeds as material for induction of mutagenesis also frequently produces chimera which single cells may be induced mutation, then divide and differentiate into parts of the plant (Acquaah, 2007). In chimeric tissues, the mutated cells are present along with normal cells which are the main bottleneck for mutagenesis induction in plant (Datta and Chakrabarty, 2009).

### Conclusion

This study emphasized the importance of colchicine treatment for mutation induction both on directly dropping on inflorescences and seed soaking. Although some characteristics such as stomatal density and guard cell size were introduced for determining the ploidy chromosome set, but in this study, only guard cell size (width and length) showed the value decreasing when colchicine concentration was increased. For the study of DNA content by the flow cytometry, except to normal diploidy (2x), many types of mixoploid occurred in both experiments. Directly treated the inflorescences with colchicine showed mixoploidy of trisomic ( $2x + 1$ ) and tetrasomic ( $2x + 2$ ) with disomic. Soaking seeds with colchicine showed many types of mixoploid with disomic (2x) such as trisomic, tetrasomic, pentasomic ( $2x + 3$ ) and hexasomic ( $2x + 4$ ). Extra chromosome number of incomplete aneuploidy and the complex of mixoploid increased when the higher concentration of colchicine was treated.

### Acknowledgements

We gratefully thank both Silpakorn University Research and Development Institute (SURDI), Thailand for funding and Faculty of Animal Sciences

and Agricultural Technology, Silpakorn University, Phetchaburi IT campus, Cha-Am, Phetchaburi, Thailand for providing laboratory and field supports during the course of this research.

### References

- Acquaah, G. (2007) *Principles of plant genetics and breeding*, Blackwell Publishing Ltd., Oxford.
- Ambrosi, D. G., Galla, G., Purelli, M., Barbi, T., Fabbri, A., Lucretti, S., Sharbel, T. F., and Barcaccia, G. (2010) DNA marker and FCSS analyses shed light on the genetic diversity and reproductive strategy of *Jatropha curcas* L. *Diversity* 2: 810-836.
- Basha, S. D. and Sujatha, M. (2007) Inter and intra-population variability of *Jatropha curcas* (L.). Characterized by RAPD and ISSR markers and development of population-specific SCAR markers. *Euphytica* 156: 375-386.
- Bever, J. D. and Felber, F. (1991) The theoretical genetics of autopolyploidy. *Oxford Surveys in Evolutionary Biology* 7: 185-217.
- Bhattacharya, A., Datta, K., and Kumar, S. D. (2005) Floral biology, floral resource constraints and pollination limitation in *Jatropha curcas* L. *Pakistan Journal of Biological Sciences* 8: 456-460. DOI: 10.3923/pjbs.2005.456.460
- Birchler, J. A. (2010) Reflections on studies of gene expression in aneuploids. *Biochemical Journal* 426: 119-123.
- Birchler, J. A., Bhadra, U., Bhadra, M. P., and Auger, D. L. (2001) Dosage-dependent gene regulation in multicellular eukaryotes: implications for dosage compensation, aneuploidy syndromes, and quantitative traits. *Developmental Biology* 234: 275-288.
- Carretero, C. L., Cantos, M., and Garcia, J. L. (2007) In vitro-ex vitro salt (NaCl) tolerance of cassava (*Manihot esculenta* Crantz)

- plants. *In Vitro Cellular and Developmental Biology-Plant* 43: 364-369.
- Dasgupta, M., Sahoo, M. R., Kole, P. C., and Mukherjee, A. (2008) Evaluation of orange-fleshed sweet potato (*Ipomoea batatas* L.) genotypes for salt tolerance through shoot apex culture under in vitro NaCl mediated salinity stress conditions. *Plant Cell, Tissue and Organ Culture* 94: 161-170.
- Datta, S. K. and Chakrabarty, D. (2009) Management of chimera and in vitro mutagenesis for development of new flower color/shape and chlorophyll variegated mutants in Chrysanthemum. In *Inducted plant mutations in the genomics era* (Shu, Q. Y., ed.), pp. 303-305. Food and agriculture organization of the United Nation, Rome.
- Doležel, J. and Bartoš, J. (2005) Plant DNA flow cytometry and estimation of nuclear genome size. *Annual of Botany* 95: 99-110.
- Fernandez, A. O. and Muzica, B. (1973) Effects of some environmental factors on the differentiation of stomata in *Spirodela intermedia* W. Koch. *Botanical Gazette* 134: 117-121.
- Ginting, C. and Maryono, T. (2009) Physic nut (*Jatropha curcas* L.) diseases in Lampung province. *Biotropia* 16: 45-54.
- Heller, J. (1996) Physic nut, *Jatropha curcas* L. *Promoting the conservation and use of underutilized and neglected crops. No 1*. International Plant Genetics Resource Institute, Rome. 66 pages.
- Henry, I. S., Dilkes, B. P., Miller, E. S., Burkart-Waco, D., and Comai, L. (2010) Phenotypic consequences of aneuploidy in *Arabidopsis thaliana*. *Genetics* 186: 1231-1245. DOI: 10.1534/genetics.110.121079.
- Heslop-Harrison, J. S. (1995) Flow cytometry and genome analysis. *Probe* 5: 14-17.
- Hill, H. D. and Myers, W. M. (1944) Isolation of diploid and tetraploid clones from mixoploid plants of ryegrass (*Lolium Perenne* L.), produced by treatment of germinating seeds with colchicine. *Journal of Heredity* 35: 359-361.
- Huang, H. -P., Gao, S. -L., Chen, L. -L., and Wei, K.-H. (2010) *In vitro* tetraploid induction and generation of tetraploids from mixoploids in *Dioscorea zingiberensis*. *Pharmacognosy Magazine* 6: 51-56.
- Inagaki, M. N. (2003) Double haploid production in wheat through wide hybridization. In *Doubled haploid production in crop plants* (Maluszynski, M., Kasha, K. J., Forster, B. P., and Szarejko, I. (eds.), pp. 53-58. Springer Science + Business Media, New York.
- Jenczewski, E., Eber, F., Manzanares-Dauleux, M. J., and Chevre, A. M. (2002) A strict diploidy-like pairing regime is associated with tetrasomic segregation in induced autotetraploids of kala. *Plant Breeding* 121: 177-179.
- Jongschaap, R. E. E., Corre, W. J., Bindraban, P. S., and Brandenburg, W. A. (2007) Claims and facts on *Jatropha curcas* L.: Global *Jatropha curcas* evolution. *Breeding and propagation programme. Research Report*. [Online URL: <http://library.wur.nl/WebQuery/wurpubs/lang/358549>] accessed on January 3, 2012.
- Koh, H. -J., Heh, M. -H., and Lee, G. -S. (1993) Transmission of extra chromosomes and its effects on anther culture in rice trisomic plants. In *Biotechnology in Agriculture* (You, C., Chen, Z., and Ding, Y. (eds.). *Current Plant Science and Biotechnology in Agriculture* 15: 313-316.
- Kung, C. Y. and Tam, H. C. (2005) On-line management and control of distributed renewable energy power plant part I-Fuzzy

- modeling of PV sytem. *PVSEC-15*, Shanghai China.
- Lee, E. A., Darrah, L. L., and Coe, E. H. (1996) Dosage effects on morphological and quantitative traits in maize aneuploids. *Genome* 39: 898-908.
- Li, M. R., Li, H. Q., and Wu, G. J. (2006) Study on factors influencing Agrobacterium-mediated transformation of *Jatropha curcas* (in Chinese). *Journal of Molecular Cell Biology* 39: 83-89.
- Mayo, O. (1971) Rates of change in gene frequency in tetrasomic organisms. *Genetica* 42: 329-339.
- Miller, M., Zhang, C., and Chen, Z. J. (2012) Ploidy and hybridity effects on growth vigor and gene expression in *Arabidopsis thaliana* hybrids and their parents. *G3 (Bethesda)* 2: 505-513.
- Mishra, M. K. (1997) Stomatal characteristics at different ploidy levels in *Coffea* L. *Annals of Botany* 80: 689-692.
- Moody, M. E., Mueller, L. D., and Soltis, D. E. (1993) Genetic variation and random drift in autotetraploid populations. *Genetics* 134: 649-657.
- Oates, K. M., Touchell, D. H., and Ranney, T. G. (2013). Induced variation in tetraploid *Rudbeckia subtomentosa* 'Henry Eilers' regenerated from gamma-irradiated callus. *HorstScience* 48: 831-834.
- Omezzine, F., Ladhari, A., Nefzi, F., Harrath, R., Aouni, M., and Haouala, R. (2012) Induction and flow cytometry identification of mixoploidy through colchicine treatment of *Trigonella foenum-graecum* L. *African Journal of Biotechnology* 11: 16434-16442.
- Paepatung, N., Nopharatana, A., and Songkasiri, W. (2009) Bio-methane potential of biological solid materials and agricultural wastes. *Asian Journal on Energy and Environment* 10: 19-27.
- Peerapong, P. and Limmeechokchai, B. (2009) Exergetic and thermoeconomic analyses of the rice-husk power plant in Thailand. *Journal of Metals, Materials and Minerals* 19: 9-14.
- Piromya, R. and Kermanee, P. (2013) Occurrence of tetraploidy in cochicine-treated physic nut (*Jatropha curcas* Linn.). *Kasetsart Journal (Nat. Sci.)* 47: 23-29.
- Rajore, S. and Batra, A. (2005) Efficient plant regeneration via shoot tip explant in *Jatropha curcas*. *Journal of Plant Biochemistry and Biotechnology* 14: 73-75.
- Sapra, V. T., Hughes, J. L., and Sharma, G. C. (1975) Frequency, size and distribution of stomata in triticale leaves. *Crop Science* 15: 356-358.
- Sardana, J., Batra, A., and Ali, D. J. (2000) An expeditious method for regeneration of somatic embryos in *Jatropha curcas* L. *Phytomorphology* 50: 239-242.
- Sharma, H. K., Shukla, A., Kumar, A., Shukla, A., Choudhary, S. B., and Jatothu, J. L. (2013) Variability and genetic diversity assessment in physic nut (*Jatropha curcas* L.). *Journal of Medical Plants Research* 7: 2380-2391.
- Songsri, P., Suriharn, B., Sanitchon, J., Srisawangwong, S., and Kesmala, T. (2011) Effects of gamma radiation on germination and growth characteristics of physic nut (*Jatropha curcas* L.). *Journal of Biological Sciences* 11: 268-274.
- Sujatha, M., Makkar, H. P. S., and Becker, K. (2006) Shoot bud proliferation from axillary nodes and leaf sections of non-toxic *Jatropha curcas* L. *Plant Growth Regulation* 47: 83-90.
- Sujatha, M. and Mukta, N. (1996) Morphogenesis and plant regeneration from tissue cultures of *Jatropha curcas*. *Plant Cell, Tissue and*

- Organ Culture (Historical Archive)* 44: 135-141.
- Sujatha, M. and Prabhakaran, A. J. (2003) New ornamental *Jatropha* hybrids through interspecific hybridization. *Genetic Resources and Crop Evolution* 50: 75-82.
- Sujatha, M. and Sailaja, M. (2005) Stable genetic transformation of castor (*Ricinus communis* L.) via *Agrobacterium tumefaciens*-mediate gene transfer using embryo exes from mature seeds. *Plant Cell Report* 23: 803-810.
- Tan, G. Y. and Dunn, G. M. (1973) Relationship of stomatal length and frequency and pollen grain diameter to ploidy level in *Bromus inermis* leys. *Crop Science* 13: 332-334.
- Trabi, M., Gübitz, G. M., Steiner, W., and Foidl, N. (1997) Toxicity of *Jatropha curcas* seeds. *In: Proceeding "Jatropha 97": Biofuels and industrial products from Jatropha curcas.* Gübitz, G. M., Mittelbach, M., and Trabi, M. (Eds), Managua, Nicaragua, 23-27 February.
- Wei, Q., Lu, W. -D. Liao, Y., Pan, S. L. Xu, Y., Tang, L., and Chen, F. (2004) Plant regeneration from epicotyl explant of *Jatropha curcas*. *Journal of Plant Physiology and Molecular Biology* 30: 475-478.
- Winkler, E., Gübitz, G. M., Foidl, N., Staubmann, R., and Steiner, W. (1997) Use of enzymes for oil extraction from *Jatropha curcas* seeds. *In: Proceeding "Jatropha curcas"* Gübitz GM, Mittelbach M, Trabi M (eds), Managua, Nicaragua, 23-27 February.